

# PRV

PATENT- OCH REGISTRERINGSVERKET  
Patentavdelningen

REC'D 23 AUG 2000

WIPO

PCT

Intyg  
Certificat



Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

This is to certify that the annexed is a true copy of ~~the documents as originally filed with the Patent- and~~ Registration Office in connection with the following patent application.

SE 00/01484

(71) Sökande A+Science Invest AB, Göteborg SE  
Applicant (s)

(21) Patentansökningsnummer 9902817-7  
Patent application number

4

(86) Ingivningsdatum 1999-07-30  
Date of filing

Stockholm, 2000-08-15

För Patent- och registreringsverket  
For the Patent- and Registration Office

Åsa Dahlberg  
Åsa Dahlberg

Avgift  
Fee

**PRIORITY  
DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

PATENT- OCH  
REGISTRERINGSVERKET  
SWEDEN

Postadress/Adress  
Box 5055  
S-102 42 STOCKHOLM

Telefon/Phone  
+46 8 782 25 00  
Vx 08-782 25 00

Telex  
17978  
PATOREG S

Telefax  
+46 8 666 02 86  
08-666 02 86

AWAPATENT AB

A\* SCIENCE INVEST AB

Kontor/Handläggare

Ansökningsnr

Vår referens

Göteborg/Maria Stenbäck

SE-2996183

1

A METHOD FOR SELECTIVE ELECTROFUSION OF AT LEAST TWO  
FUSION PARTNERS HAVING CELL-LIKE MEMBRANES

Field of the invention

The present invention relates to a method for selective electrofusion of at least two fusion partners having cell-like membranes.

5

Background of the invention

Electrofusion has developed into an extremely efficient method for the fusion of mammalian cells, mainly because of its mild conditions, which result in a high number of viable fusion products [see e.g. White, K. L. 1995, Electrofusion of mammalian cells, Methods in Molecular Biology, 48, 283-293]. The application of an electrical field over phospholipid bilayer membranes induces pore formation when the applied potential reaches or exceeds the membrane breakdown potential. Consequently, electro-permeabilization techniques has been used in a wide variety of biological experiments, like electrofusion for the creation of hybridomas and new cell lines [see e.g. Zimmermann, U., et al., 1985, Electrofusion: a novel hybridization technique, Adv. Biotechnol. Proc. 4, 79-150; Neil, G. A. et al., 1993. Electrofusion, Methods in Enzymology, 220, 174-196; Glassy, M. 1988, Product review: Creating hybridomas by electrofusion, Nature, 333, 579-580], in vitro fertilization [see e.g. Ogura, A. et al., 1995, Spermatids as male gametes. Reprod. Fertil. Dev., 7, 155-159], cloning experiments [see e.g. Van Stekelenburg-Hamers, A. E. P., et al., 1993, Nuclear transfer and electrofusion in bovine in vitro-matured / in vitro-fertilized embryos: effect of media and electrical fusion parameters, Mol. Reprod. Dev., 36, 307-312], electroporation of cells for introduction of cell-impermeant solutes [see e.g. Electroporation: a gen-

eral phenomenon for manipulating cells and tissues, J. Cell. Biochem., 51: 426-435; Li, H., et al., 1997, Transfection of rat brain cells by electroporation, J. Neurosci. Methods, 75, 29-32; Lundqvist, J. A., et al., 1998, Altering the biochemical state of individual cultured cells and organelles with ultramicroelectrodes, Proc. Natl. Acad. Sci. USA, 95, 10356-10360], and electroinser-

---

tion for the addition of membrane-associated macromolecules, including proteins [see e.g. Mouneimne, Y., et al., 1989, Electro-insertion of xeno-glycophorin into the red blood cell membrane, Biochem. Biophys. Res. Com. 159, 34-40]. Applications of *in vivo* electrofusion include the incorporation of gonococcal attachment receptors from human HL60 cells to rabbit corneal epithelial tissue as a viable model of human-specific pathogens [see e.g. Heller, R., et al., 1990, Transfer of human membrane surface components by incorporating human cells into intact animal tissue by cell-tissue electrofusion *in vivo*, Biochim. Biophys. Acta, 1024, 185-188].

Electric-field-induced fusion is widely employed in biomedical research for a population of cells in suspension. Cells are first brought into contact by dielectrophoresis through the application of a low-amplitude, high-frequency AC field and subsequently a fraction of the cells are fused by a strong and short DC pulse. Bulk electrofusion of large quantities of cells is useful for creating and selecting new cell lines, but cannot be applied to fuse single cells with high precision. This leads to unwanted fusion between cells from the same cell-line, as well as the wanted fusion between cells from different cell lines. Furthermore, bulk electrofusions do not allow the control over the number of cells that are to be fused together, which leads to unfavorable ratios of dinuclear-to-multinuclear fusion products.

# Summary of the invention

It is often desirable to selectively and controllably alter the biochemical, and genetic properties of single cells. To address this challenge and to overcome the shortcomings of bulk electrofusion, the inventors of the present invention have developed a technique to fuse together a single pair of cells at a time. The ability to controllably fuse together single cells represents a technique by which the long-term genetic identity and behavior of a selected cell can be precisely manipulated, and opens up new possibilities to create combinatorial libraries of hybrid, and cloned cells. In combination with a powerful measurement and imaging technique, the genetic and biochemical nature of single cells can be studied in detail.

Alteration of cellular properties can also be achieved by fusing together a synthetic phospholipid vesicle with the desired vesicular content and membrane compositions to a target cell. This technique can be used to alter the contents and membrane properties of single cells. It is, for example, possible to introduce a membrane protein reconstituted in liposomes into the cell plasma membrane. This ability to selectively transform the membrane composition of single cells is anticipated to have useful biological applications, such as the introduction of surface receptors for the screening of potential ligands and related pharmacological compounds.

Thus, the present invention provides a novel method for the selective electrofusion of cellular structures, such as cells and liposomes. This method offers the advantage of cell-selection, fusion of adherent cell structures with a high spatial resolution, giving the possibility to create complex cellular networks. As electrofusion is a mild method, this miniaturized version can, for example, be used for cloning on the single cell level and for *in vitro* fertilization. For cloning experiments in particular, the shortcomings of bulk electrofusion is

overcome with the present technique as it offers complete control over the fusion process and any doubts to the identity of the somatic cells needs not to be raised. Also, this method, preferentially in combination with micro-  
 5 machined chip technology, can be used to create screening libraries of cloned cells or hybrid cells.

The characterizing features of the invention will be evident from the following description and the appended claims.

10

Brief description of the drawings

In the description and the examples below, reference is made to the accompanying drawings on which:

Fig. 1 is a schematic drawing showing the experimental set up. Cells 1, 2 are added from suspension to a microscope coverslip 3 mounted in a polycarbonate holder. The cells are either prealigned using optical trapping (MOPA laser 4) or simply by pushing them together with the microelectrodes 5, 6 controlled by high-graduation  
 15 micromanipulators 7. For fluorescence imaging and optical trapping, two collinear laser beams are sent into the microscope objective 9; an argon ion laser 10 (black, thin arrows) is used to excite fluorescein, and a MOPA diode laser 4 (black thick arrows) is used for optical trap-  
 20 ping. The resulting fluorescence and bright field images are directed to a CCD camera 11.

Figs. 2 A - C show the alignment of a liposome with a selected cell using a microelectrode, and Figs. 2 D - E show six fluorescein-containing (10  $\mu$ M) liposomes brought  
 30 into contact with a cell. As shown in Fig. 2 G, no resulting background fluorescence is detected, and this illustrates that selective delivery of a drug to the surface (electropermeabilization of the liposome), or interior (electrofusion between the cell and liposome) is  
 35 possible.

Figs. 3 A - D show a cell-cell fusion sequence. Fig. 3 A shows PC12 cells during dielectrophoresis (0.3 - 3

kV/cm, 2 MHz). Fig. 3 B illustrates initiation of the fusion by the last fusion pulse (6 pulses of 1 ms duration per each, 3 kV/cm), this can be seen by a broader and flatter contact zone between the cells. Fig. 3 C (~1 minute after the fusion pulses) and Fig. 3 D (~2 minutes after the fusion pulses) show the subsequent broadening of the fusion area between the cells, indicating complete fusion.

---

Figs. 4 A - C show Bright-field images taken before (Fig. 4 A), during (Fig. 4 B), and after (Fig. 4 C) electrofusion (8 kV/cm, 4 ms) of a phosphatidylcholine (PC) liposome (left) with a COS 7 cell (right). Figs. 4 D - E show fusion between a NG-108 cell (protease-treated for 30 minutes) and a PC liposome with incorporated  $\gamma$ -glutamyltransferase ( $\gamma$ -GT). In Fig. 4 E it is seen how the microelectrode is gently pulled away from the cell, and the liposome (which is attached to the microelectrode) is stretched but does not detach from the cell.

Fig. 5 illustrates an example of a clinical application of the method according to the invention, wherein single or multiple cells in the brain are electrofused to incorporate the content of the other fusion partner. The fusion partner is delivered via a hollow electrode. The electrode is positioned using a stereotactic device, represented in the figure by the Cartesian co-ordinate system, and stereotactic micropositioners.

#### Detailed description of the invention

The invention relates to a method for selective electrofusion of at least two fusion partners having cell-like membranes, comprising the following steps: A) the fusion partners are brought into contact with each other, and B) an electrical field of a strength sufficient to obtain fusion and highly focused on the fusion partners is applied.

Said at least two fusion partners having cell-like membranes are preferably structures of cellular or sub-

cellular dimensions. The expression "structure of cellular or subcellular dimensions" relates in particular to biological structures such as independent cells and smaller structures, but it also relates to similar artificial structures. Thus, the fusion partners may be, independently of each other, a single cell, a liposome, a proteoliposome, a synthetic vesicle, a plant protoplast, an egg cell, a sperm or spermatid, and an enucleated egg cell.

As stated above, the fusion partners are brought into contact with each other. This means either that they are placed so that the outer surfaces of the fusion partners are touching, or that the fusion partners are placed at a very small distance from each other.

In most cases the method according to the invention is used for fusion of two fusion partners, below denoted Fusion Partner I and Fusion Partner II, however it, is also possible to fuse more than one fusion partner. In some cases it is especially interesting to fuse more than two partners, for example, to create multinuclear cells with more than two nuclei, or when one of the partners is of much larger dimensions than the others. An example of this is when one wishes to introduce a substance contained in vesicles into a cell. In this case Fusion Partner I is the cell and Fusion Partner II is several of the smaller vesicles.

Furthermore, it is possible to repeat the method according to the invention one or several times, so that a new fusion partner is fused to two already fused fusion partners. This is of particular importance when combinatorial libraries of cloned cells and hybrid cells are created.

When Fusion Partner I is a cell and Fusion Partner II is a single liposome, the fusion of them allows both the introduction of the liposomal content into the cell interior as well as the addition of lipids and membrane proteins from the liposome membrane into the cell sur-

face. This cell-liposome fusion represents a novel approach to the manipulation of the membrane contents and surface properties of single cells.

To fuse Fusion Partner I, such as a cell, a liposome  
5 or another similar structure, to Fusion Partner II, such  
as a cell, liposome or similar structure, it is necessary  
to place the two fusion partners next to each other, i.e.  
~~in contact with each other. The fact that the two fusion~~  
partners are placed next to each other before fusion  
10 makes it possible to avoid dielectrophoresis, which tra-  
ditionally is used for the creation of close contact be-  
tween cells. However, dielectrophoresis can be used suc-  
cessfully in combination with this invention to create  
and establish close cell-cell contacts before the dc-  
15 field fusion pulse is applied. Mechanically, close con-  
tact between the fusion partners can be done in any suit-  
able way. Individual manipulation of the two fusion part-  
ners for alignment facilitates this alignment. Using, for  
example, optical trapping with highly focused laser  
20 beams, individual cells, as well as other biological  
structures, including organelles of small dimension, can  
be manipulated and moved at will [this has been described  
previously, see e.g. Jaroszeski, M. J., et al., 1994. Me-  
chanically facilitated cell-cell electrofusion, Biophys.  
25 J., 67, 1574-1581; Uchida, M., et al., 1995, Whole-cell  
manipulation by optical trapping, Curr. Biol., 5, 380-  
382; Chiu, D. T., et al., 1998, Probing single secretory  
vesicles with capillary electrophoresis, Science, 279:  
1190-1193]. When electrodes are used for positioning of  
30 the fusion partners, as described below, it is possible  
to align the two fusion partners by adjusting their posi-  
tions by moving them with the tips of the electrodes. In  
order to move the electrodes it may be advantageous to  
use a microscope, at least one micropositioner and/or a  
35 stereotactic device. To further facilitate the position-  
ing of the two fusion partners, it may be advantageous to  
immobilize one of the fusion partners prior to performing



step A. When electrodes are used, this may, for example, be done by reversibly adhere at least one of the fusion partners to at least one of the microelectrodes. In some cases it may be advantageous to use dielectrophoresis, as pretreatment of the two aligned cells, before fusion. This can be done by connecting an ac-field sweep function generator to the electrodes.

The electrical field may be obtained by use of a low-voltage or high-voltage pulse-generator depending on the electrodes used, and other experimental parameters. The voltage generator is used to produce an electric field strength sufficient to result in fusion between the two fusion partners, approximately 0.1-10 kV/cm, for durations of 10  $\mu$ s to several seconds. The voltage measured at the membrane of the fusion partners should be between a few hundred millivolts to several volts, preferably around 1.5V. In the case of multiple-voltage-pulse protocols, we have found that a pulse repetition rate of approximately 1 Hz, to be suitable, however, other repetition rates might work well. For longer pulse-application durations, correspondingly, repetition rates of a lower frequency should be used. In any event, the length and the strength of the pulses depend on the size of the partners to be fused. Preferably, the fusion pulses have a rectangular waveform, but other waveforms works as well, including various ac-field pulse protocols. For dielectrophoresis, the sweep function generator preferably generates a alternating field, sinus-wave form, of field strengths between 100 V/cm - 5 kV/cm, 100 Hz - 2 MHz.

The electrical field used in step B to obtain fusion should be highly focused in order to avoid affecting any surrounding structures and to obtain the advantages of the present invention. To focus the electrical field it is preferable to provide the electrical field by use of one or two microelectrodes positioned close to the two fusion partners, i.e. 0 - 10  $\mu$ m, preferably 0 - 5  $\mu$ m, from the cellular membrane. In the case of using a single

electrode, this electrode is preferably, biased at a positive potential (anode) and work against a grounded cell preparation. According to the invention the micro-electrodes are preferably electrodes of cellular to sub-cellular dimensions. Preferably the outer dimension of the ends of the electrodes positioned closest to the fusion partners is from a few nanometers to ~100 micrometers, more preferably 5-30 micrometers and most preferably approximately 20  $\mu\text{m}$ . The electrodes can be made of a solid electrically conducting material, or they can be hollow for delivery of fusion partners or chemical agents of choice. The electrodes can be made from different materials. A special type of electrodes are hollow and made from fused silica capillaries of a type that frequently is used for capillary electrophoresis and gas chromatographic separations. These capillaries are typically one-to-one hundred micrometers in inner diameter, and five-to-four hundred micrometers in outer diameter, with lengths between a few millimeters up to one meter. For cell fusion applications, these electrodes are filled with an electrolyte, preferably a physiological buffer solution. When a potential sufficient to cause cell-fusion is applied over the capillary, electroendoosmotic bulk flow is induced in the capillary, which in combination with Pouiseille flow (gravitational flow in capillaries) can be used to efficiently transfer materials to the fusion partners. Such hollow narrow-bore fused silica capillaries have the additional advantage that components added to the inlet end can be fractionated based on their charge-to-frictional drag ratio. This characteristic feature of the system can be used to, for example, investigate the effect of various fractionated components on cell fusion.

Furthermore, it is preferred to provide the two fusion partners in an electrofusion buffer prior to step B.

In order to facilitate the fusion, it may be advantageous to pre-electroporate at least one of the fusion partners before step A is performed.

When one (or both) of the fusion partners is a cell, it may be part of a cellular network or a tissue, for example, in order to study cellular networks where, for example, the alteration of the biochemical identity of a selected cell can have profound effects on the behavior of the entire network.

A setup suitable for performing the method according to the invention is illustrated in Fig. 1. In this figure the two fusion partners 1, 2 both are cells. The cells to be fused are added from a suspension to a microscope coverslip 3 mounted in a polycarbonate holder.

The cells are either prealigned using optical trapping (MOPA laser 4) or simply by pushing them together with the microelectrodes 5, 6 controlled by high-graduation micromanipulators 7. The microelectrodes are preferably carbon fiber electrodes, and most preferably carbon fiber ultramicroelectrodes (5  $\mu\text{m}$  in diameter), or hollow glass fiber electrodes.

A voltage generator 8 is used to provide the required electrical field.

For fluorescence imaging and optical trapping, two colinear laser beams are sent into the microscope objective 9. A 488-nm beam from an argon ion laser 10 (the beam from the argon ion laser is denoted by black, thin arrows) was used to excite fluorescein and a 992-nm beam from a MOPA diode laser 4 (the beam from the argon ion laser is denoted by black, thick arrows) was used for optical trapping.

The resulting fluorescence and bright field images are directed to a CCD camera 11.

The setup also comprises mirrors 12, a dichroic beamsplitter 13, a polychroic beamsplitter 14, a lens 15, a spinning disc 16, and a filter 17.

The method according to the invention is suitable for manipulating the genetic identity and biochemical surface properties of individual cellular or sub-cellular structures, such as organelles. It is especially suitable and interesting for in vitro-fertilization. It can also be used in several other applications such as cloning, creation of hybridomas, manipulation of the composition of a cellular membrane, and delivery of a well-defined volume of a substance to a cell (particularly delivery of a pharmaceutically active substance to a cell).

The method according to the invention is also suitable for electrofusion in vivo of patients or animals suffering from biochemical disruptions in individual groups of cells.

Many diseases, may they be genetically acquired or not, results in metabolic disruptions. For example, Parkinson's disease caused by degeneration of neurons in the Nigro-striatal pathway result in malfunctioning in the biochemical machinery for production of dopamine in an isolated population of cells. This in turn results in motorbehavioral deficits. The standard treatment of Parkinson's disease is by oral administration of L-DOPA, a precursor of dopamine. Alternatively, grafted tissue with neuronal cells producing dopamine is transplanted into the patient's brain. Intracellular drug or gene-administration in vivo into the appropriate brain structures can be accomplished using an electrofusion procedure similar to that described in the above examples and used as a therapeutic strategy.

Other diseases where a small population of cells are malfunctioning include many disorders in the inner organs, but also tumors.

Experimental treatment of brain tumors using genetically engineered viruses for gene delivery has been used with anecdotal reports of success. However, the use of viruses as a delivery system has limitations in that it might pose a potential hazard should the virus mutate. Using an electrofu-

sion procedure for gene delivery (e.g. "suicide genes" cytokine deaminase or thymidine kinase) similar to that described in the examples below eliminates the need for the use of virus delivery systems in cancer therapy.

- 5 Many times a focal administration (administration directly to the malfunctioning set of cells) of drugs or genes (gene therapy) can be expected to be far superior than intraperitoneal, oral, intraventricular, or any other kind of commonly employed drug-administration technique. Intracellu-
- 10 lar drug-and-gene-administration in vivo can be accomplished using an electrofusion procedure similar to that described in the examples below. Because of the extremely small dimensions of the electrodes, in combination with the low voltages applied, very little tissue trauma is expected. Furthermore,
- 15 the positioning of the electrodes and the subsequent gene or drug delivery is very precise. This is especially important in the brain. It has been shown that microdialysis probes, which are on the order of a hundred times larger than the electrodes employed here for electroporation, cause very
- 20 little tissue trauma and disruption of local metabolism 24 hours after implantation.

For solute transfer into cells and organelles in vivo using electroporation it is preferred to use electrodes which are hollow. The fusion partner to be electrofused into cells

25 are then simply administered through the narrow channel in the center of the electrode by application of a flow by means of a syringe pump or a peristaltic pump or any other type of solution pumping system including electrophoresis.

An especially interesting possibility is to use battery-

30 operated perfusion/electroporation implants in bio-tolerable materials for continuous application of solutes with fusion partner for electrofusion into cells or tissue. Because such low potentials are required, batteries with emf's in the range of a few to 20 volts can be used. These battery-

35 operated electroporation units can be made small, virtually they can be included on a chip measuring only a few millimeters squared.

An example of an *in vivo* setup is shown in Fig. 5. A hollow electrode 20 is positioned by use of a stereotactic device close to a cell 21 in the brain. This cell 21 constitutes one of the fusion partners. The other fusion partner is  
 5 constituted by liposomes containing a pharmaceutically active substance, such as DNA, a protein, RNA or a drug. The cell is perfused with the liposomes by injection of the second fusion partner into the electrode 21, ~~either by use of a syringe 22~~  
 as shown in the figure, or by use some other appropriate  
 10 means.

The invention will now be further explained in the following example. This example is only intended to illustrate the invention and should in no way be considered to limit the scope of the invention.

15

#### Examples

In the examples, the following methods and materials were used.

#### 20 **Micromanipulation and fusion**

Optical trapping with highly focused laser beams was used to align the fusion partners. Using this procedure, one of the fusion partners (cell or liposome) in solution was trapped by an IR laser beam and brought into contact  
 25 with the other fusion partner, which was either an adherent cell grown on a substratum or a cell placed on the substratum using, optical trapping with the set-up schematically shown in Fig. 1, and described above.

For positioning of cells or liposomes close to cellular  
 30 targets for fusion, the inventors also use reversible adsorption to carbon fiber microelectrode tips. In using this particular scheme for cell-liposome fusions, liposomes co-immobilized with cells on poly-L-lysine-coated glass surfaces are detached from the surface by pushing  
 35 the electrode tip against the liposome roughly parallel to the object plane. The liposome will then adhere lightly to the electrode tip and can be relocated close

to a cellular structure by micromanipulators that can be moved in increments of 0.2  $\mu\text{m}$  in three dimensions. This is shown in Figs. 2 A - C, where a liposome of about 5  $\mu\text{m}$  in diameter is moved about 90  $\mu\text{m}$  to a target cell, while  
5 Figs. 2 D - G show how complex cell-liposomal patterns can be created using this simple approach. The liposomes in Figs. 2 A - C were first immobilized on poly-L-lysine-coated boro-silicate coverslips, and cells were added in suspension. A liposome could easily be moved over 90  $\mu\text{m}$   
10 to a cell. One advantage of this technique over optical trapping is the much larger force that can be applied to the vesicles. The same method can be used for the alignment of two cells, with the exception that poly-L-lysine-coated coverglasses do not have to be used as the cells  
15 in suspension simply attach to the surface due to gravity.

For electrofusion, the same pair of 5  $\mu\text{m}$ -outer diameter carbon fiber microelectrodes is used. Fig. 1 depicts the geometrical arrangement of the electrodes with  
20 respect to a pair of cells prealigned for fusion. For cell-cell, and cell-liposome fusion, highly focused electrical fields are applied using a low-voltage pulse-generator. An advantage of this set-up is that the electrodes are of cellular to subcellular dimensions, enabling  
25 fusion of single cells in complex cellular networks grown on a substratum. At the same time, the highly focused electric field minimizes the risk for unwanted fusion or electroporation of surrounding cells.

### 30 **Experimental protocol**

#### **Optical trapping, microscopy, and fluorescence imaging**

The optical trapping and fluorescence imaging systems were built in-house. The optical trap was formed by sending the output from a single-mode MOPA laser diode  
35 (Model SDL-5762-A6, SDL, Inc.) through a spatial filter (model 900, Newport) so that the output was then reflected off a near-IR mirror. The reflected laser light

was passed through a dichroic mirror, reflected from a polychroic mirror (Chroma Technology Corp.) that is placed in a microscope (Nikon Diaphot or Leica DM IRB), and subsequently brought to a diffraction-limited focus with a high-numerical aperture objective (100 x, NA 1.4 Nikon and 100 x, NA 1.3 Leica, respectively). Fluorescence excitation was achieved by sending the 488-nm output of an argon ion laser (Spectra Physics 2025-05) through a telescope followed by a spinning disc. The purpose of the spinning disc is to scatter the laser light so that uniform illumination is achieved for fluorescence imaging. The scattered laser light from the disk was collected by a lens and reflected from a dichroic mirror (Chroma Technology Corp). This reflected light was sent into the microscope and was reflected by a polychroic mirror. Fluorescence and brightfield imaging was performed by a 3-chip color CCD-camera (Hamamatsu) or a silicon-intensified target camera (Hamamatsu) and recorded by a Super VHS recorder (Panasonic) or dumped directly to a hard disk via a frame grabber.

#### **Electrofusion instrumentation**

For electrofusion experiments, cell dishes were mounted in a circular polycarbonate holder using vacuum grease, and transferred to the stage of an inverted microscope. Two carbon fiber microelectrodes (Axon Instruments, Inc. ProCFE) with an outer diameter of 5  $\mu\text{m}$  were positioned close at each side of the respective fusion partner by high-graduation micromanipulators (Narishige MWH-3). The two carbon fiber electrode tips (anode and cathode) were positioned at an angle of 0-20°, and 160-180° with respect to the object plane. Cells were fused with multiple 1-ms pulses, using a pulse generator (Digitimer Stimulator DS9A) or a homebuilt variant. The voltage output from the pulse generator was calibrated using high-impedance electrodes and a patch-clamp amplifier.



The electrofusion buffer was either a 0.3 mol/kg Hanks Hepes solution (137 mM NaCl, 5.4 mM KCl, 0.41 mM MgSO<sub>4</sub>, 0.4 mM MgCl<sub>2</sub>, 1.26 mM CaCl<sub>2</sub>, 0.64 mM KH<sub>2</sub>PO<sub>4</sub>, 3.0 mM NaHCO<sub>3</sub>, 5.5 mM D-glucose, 20 mM Hepes, pH adjusted to 7.4 with NaOH) or a 0.2 mol/kg Hepes Saline buffer (135 mM NaCl, 5 mM KCl, 10 mM glucose, 2 mM MgCl<sub>2</sub>, 2 mM CaCl<sub>2</sub>, and 10 mM HEPES) diluted with 30-50% MilliQ-water. For PC12 cells were standard iso-osmolar and hypo-osmolar fusion media used, 300 L3 and 75 L3 receptively [Foung, S. et al, J of Immunol Methods, 134, 35-42].

For the NG-108 and Cos 7 cell-cell fusion experiments, 5% PEG 4000 was added to the fusion media, and in the cell-liposome fusions, 1.25% DMSO was added.

### 15 Cell culture

Jurkat, NG 108, COS 7, and PC12 cells were cultured according to standard procedures. The NG-108 and Cos 7 cells were treated with protease (from Aspergillus Oryaze) 2mg/ml, for 5-30 min in incubator (37 degrees Celsius, 90% humidity and 5% CO<sub>2</sub> atmosphere).

### Preparation of fluorescein-encapsulated unilamellar vesicles and fluorescently labeled proteoliposomes

Liposomes from L- $\alpha$ -phosphatidylcholine in chloroform (powder from fresh egg yolk, Sigma St. Louis, MO) were obtained in high yield using a rotary evaporation method [as described e.g. in Moscho, A., et al., 1996, Rapid preparation of giant unilamellar vesicles, Proc. Natl. Acad. Sci. USA 93: 11443-11445]. Fluorescently labeled  $\gamma$ -GT was prepared by reacting with 2.3 mM fluorescein isothiocyanate (FITC). Unreacted FITC was removed by running the  $\gamma$ -GT-FITC solution through an Econopac 10DG column (BioRad Laboratories, CA), with a 6000 D cut-off. Labeled  $\gamma$ -GT was then incorporated into the vesicle during the vesicle formation process.

**Preparation of poly-L-lysine-coated cover slips**

Cover-slips (borosilicate, 28 mm in diameter, 0,13 - 0,17mm thick, Kebo, Sweden) were washed in 70% ethanol/water solution (v/v), followed by MilliQ-water. The  
5 glasses were placed in a poly-L-lysine (Sigma-Aldrich Europe)/MQ-water solution (0,1% (w/v)). Cover slips were mounted in a circular polycarbonate holder using vacuum  
grease. 1 ml of liposome solution was added. After 30  
minutes a sufficient number of liposomes were immobi-  
10 lized, and the outside solution could be changed and cells added.

**Micromanipulation of liposomes and cells using microelectrodes**

15 For translation of liposomes to different locations on poly-L-lysine-coated borosilicate surfaces, carbon fiber microelectrodes controlled by high-graduation micromanipulators (Narishige, Japan, 0.2  $\mu$ m resolution) were used simply by using a combined pushing/scraping movement  
20 to loosen the vesicles from the surface. After the vesicles were detached from the surface, they adhered to the electrode tip and could be moved over long distances and be launched close to a cellular target. To determine  
whether significant loss of the liposomal content is a  
25 result of this handling, liposomes were primed with fluorescein (10  $\mu$ M), attached to the cover slips and washed thoroughly so the fluorescence background was significantly reduced. Comparison between the intravesicular  
fluorescence intensity before and after moving the vesi-  
30 cles did not show signs of substantial loss of content (data not shown).

**Chemicals and materials**

Hepes (>99%), sodium chloride, potassium chloride,  
35 and sodium hydroxide (all Suprapur), calcium dichloride, magnesium dichloride, magnesium sulfate, potassium dihydrogen phosphate, PEG 4000 and sodium hydrocarbonate,

were purchased from Merck. D-glucose (AnalaR) was from BDH Limited Poole and fluorescein (GC-grade),  $\gamma$ -GT, Protease (type XXIII, from *Aspergillus oryzae*), fluorescein (sodium salt), and DMSO were obtained from Sigma-Aldrich, Sweden. Fluorescein isothiocyanate was from Molecular Probes, Europe. Deionized water from a Milli-Q system (Millipore) was used.

#### Example 1 - cell-cell fusion

10        In this example two PC12 cells ( $\approx 8$  and  $10 \mu\text{m}$  in diameter) brought into contact by optical trapping and immobilized on a cover slip were fused, as shown in Figs. 3 A - D. Fig. 3 A shows PC12 cells during dielectrophoresis (0.3 - 3 kV/cm, 2 MHz). Fig. 3 B illustrates initiation  
15        of the fusion by the last fusion pulse (6 pulses of 1 ms duration per each, 3 kV/cm), this can be seen by a broader and flatter contact zone between the cells. Fig. 3 C ( $\sim 1$  minute after the fusion pulses) and Fig. 3 D ( $\sim 2$  minutes after the fusion pulses) show the subsequent  
20        broadening of the fusion area between the cells, indicating complete fusion.

To demonstrate the high spatial resolution achieved with this technique, individual NG 108 cells in a network were fused (data not shown). In addition to PC 12 cells,  
25        and NG 108 cells, Jurkat cells, and Cos 7 cells, were successfully fused (data not shown). Hybrid cells between PC12 cells and NG 108 cells were successfully created in similar fusion protocols (data not shown).

Because of the presence of cytoskeleton, fused cells  
30        take much longer to reorganize its membrane and become spherical compared to liposomes. For example, typical liposome-liposome fusion, using an identical experimental set-up, is in the milliseconds regime (data not shown). For some cells with extensive cytoskeletal scaffolding,  
35        membrane reorganization can take minutes. However, fusion as characterized by the initial cytoplasmic continuity still occurs within a second. It was generally observed

that fusion of cells pre-electroporated in a 0.2 mol/kg buffer were much faster than the fusion of cells held in an isotonic 0.3 mol/kg buffer. The fusion process from initial cytoplasmic continuity to totally fused cells in an isotonic media took several hours. This is consistent with earlier accounts on the fusion media's impact on membrane fusion events, like exocytosis [as described e.g. in Weber, G. et al., 1992, Manipulation of cells, organelles, and genomes by laser microbeam and optical trap, *Int. Rev. Cyt.*, 133: 1-41] and cell-cell fusion [as described e.g. in Zimmerberg, J., et al., 1980, Micromolar  $\text{Ca}^{2+}$  stimulates fusion of lipid vesicles with planar bilayers containing a calcium-binding protein, *Science*, 210, 906-908; Scmitt, J. J., et al., 1989, Enhanced hybridoma production by electrofusion in strongly hypo-osmolar solutions, *Biochim. Biophys. Acta*, 983, 42-50; Scmitt, J. J., et al., 1989, Electrofusion of osmotically treated cells. *Naturwissenschaften*, 76, 122-123]. Lately it has been argued that the electrofusion enhancement reached by hypo-osmotic fusion media is due to spectrin denaturation [as described e.g. in Zimmermann, U., et al., 1990, Efficient hybridization of mouse-human cell lines by means of hypo-osmolar electrofusion, *Journal of Immunological Methods*, 134, 43-50; Chernomordik, L. V., et al., 1991, Evidence that the spectrin network and a nonosmotic force control the fusion product morphology in electrofused erythrocyte ghosts, *Biophys. J.*, 60, 1026-1037].

In all NG-108 and Cos 7 fusion experiments, the cells were treated with protease (5-30 min), and for the NG-108 cells the cell fusion media contained 5% (w/v) PEG 4000 [as described e.g. in Sowers, A. E., 1995, Membrane skeleton restraint of surface shape change during fusion of erythrocyte membranes: evidence from use of osmotic and dielectrophoretic microforces as probes, *Biophys. J.*, 69, 2507-2516]. The requirement of different pre-treatment and fusion conditions for the respective cell-

lines are in agreement with previous observations for cell fusion.

It is well-established that electrically fused cells are biologically intact and continue to grow. Also in our fusion protocol, cells adhered to surfaces and continued to grow when inspected five-to-twentyfour hours post-fusion.

---

#### Example 2 - cell-liposome fusion

10 Previous accounts of cell fusion have demonstrated that in addition to mammalian cells of other phenotypes, synthetic vesicles, as well as plant protoplasts can be fused to mammalian cells. Synthetic vesicles have been fused with cells using some kind of chemical treatment or  
15 such, for example PEG [as described, for example, in Stoicheva, N. G., et al., 1994, Electrically Induced fusion of mammalian cells in the presence of polyethylene glycol, J. Membr. Biol., 140: 177-182] or by using HVJ-glycoprotein-reconstituted liposomes [as described in  
20 Seibicke, S., et al., 1988, Fusion of lipid vesicles with ascites tumor cells and their lipid-depleted variants, Studies with radioactive- and fluorescent-labeled vesicles, Biochim. Biophys Acta, 944, 487-496]. Synthetic lipid-films have been incorporated in the plasma membrane  
25 in cells, and micro-injection facilitated, by gentle contact between a lipid-coated glass micropipette and the cellular membrane [as described e.g. in Laffafian, I., et al., 1998, Lipid-assisted microinjection: introducing material into the cytosol and membranes of small cells,  
30 Biophys. J., 75, 2558-2563]. Fusion schemes involving synthetic vesicles can be used for purposes of transferring cell-impermeant molecules into a target cell through mixing of contents and for transfer of membrane lipids and membrane associated structures, such as proteins,  
35 into the target cell membrane.

Figs. 4 A - E are two sequences of images that demonstrate single vesicle-cell fusion according to the inven-

tion. In the first sequence (Figs. 4 A - C) a  $\approx 5 \mu\text{m}$  diameter phosphatidylcholine vesicle is fused with a  $\approx 12 \mu\text{m}$  diameter Cos-7 cell, and in the second sequence (Figs. 4 D - E) a  $\approx 3 \mu\text{m}$  diameter phosphatidylcholine vesicle with reconstituted  $\gamma$ -GT is fused with a  $\approx 20 \mu\text{m}$  diameter NG-108 cell (protease-treated for 30 min). This demonstrates the ability of introducing liposome-incorporated proteins into a cellular structure. It is noted that fusion of vesicles with cells was much more difficult to accomplish than fusion between two cells of the same type. Both addition of DMSO ( $\approx 2\%$ ) to 0.3 mol/kg fusion medium and use of 0.2 mol/kg fusion media were found to facilitate fusion. DMSO has been shown to facilitate both the uptake of DNA using electroporation and increases the yield of fusion, and the use of hypo-osmolar medium is well-known to increase the fusion yield of cells (see above). In the experiment shown in the figure, the fusions were performed in a HEPES-buffered saline solution, and 1.25% dimethylsulfoxide and 20% MQ-water was added to the external buffer solution to assist vesicle-cell electrofusion.

### Example 3 - cell fusion using single open-bore fused silica capillaries

Fusion experiments were performed with NG 108 cells, cultivated according to standard procedures and plated onto no. 1 circular cover slips, mounted in a circular polycarbonate holder and transferred to an inverted microscope stage. During all experiments the cell-bath was grounded by a platinum wire. Cells were aligned for fusion using optical trapping. A high voltage power supply (Bertrand, Hicksville, NY, USA) operated at positive potentials of 1-30 kV was used together with electrolyte-filled fused silica capillaries (30 cm long, 30 mm i.d., 375 mm o.d.) employed as electrodes. The experimental setup was otherwise identical to that depicted in figure 1. The fused silica electrodes were sharpened to an approximate outer diameter of 50  $\mu\text{m}$  by grinding the tip

with a rotating sandpaper dish. The capillary and buffer container (that connects the inlet end of the capillary to the high-power voltage supply by a platinum wire) was filled with Hanks buffered saline. The sharpened capillary tip (outlet end) was positioned using high-graduation micromanipulators close to one of the fusion partners, typically, less than 10  $\mu\text{m}$ . Fusion of two aligned NG108-15 cells was performed by applying pulses of 5-to-15 kV for 0.1-5 seconds.

- 5
- 10        It was demonstrated that this method according to the invention is highly efficient for fusion of two cells in a complex system. This is especially important for application in in vivo fusion of cells and other fusion partners.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22

CLAIMS

1. A method for selective electrofusion of at least two fusion partners having cell-like membranes, comprising the following steps:

A) the fusion partners are brought into contact with each other

B) ~~an electrical field of a strength sufficient to obtain fusion and highly focused on the fusion partners is applied.~~

2. A method according to claim 1, wherein only one microelectrode, sufficiently small to permit the selective fusion of two fusion partners, are used to provide the electrical field in step B.

3. A method according to claim 1, wherein two microelectrodes, sufficiently small to permit the selective fusion of two fusion partners, are used to provide the electrical field in step B.

4. A method according to claim 1 or 2, wherein one electrode mounted on a microchip is used to provide the electrical field in step B.

5. A method according to claim 1 or 3, wherein several electrodes mounted on a microchip are used to provide the electrical field in step B.

6. A method according to claim 4 or 5, wherein the electrode(s) is (are) mounted on a microchip of a suitable design for combinatorial synthesis of fusion products.

7. A method according to any one of the claims 2-6, wherein at least one microelectrode is hollow, and sufficiently small to permit the selective fusion of two fusion partners.

8. A method according to claim 2-7, wherein at least one microelectrode that is hollow, electrolyte-filled, and sufficiently small to permit the selective fusion of two fusion partners, is used to provide the electrical field in step B, and said microelectrode is also used to



deliver fusion partners or chemical agents by electroendosmosis, electrophoresis, or by Poiseuille flow.

9. A method according to any one of the claims 2-8, wherein the outer diameter of said electrode(s) is sufficiently small to permit the selective fusion of said at least two fusion partners without affecting nearby structures, such as cells, liposomes, and proteoliposomes).

~~10. A method according to claim 9, wherein the outer diameter of said electrode(s) is 1-100  $\mu$ m.~~

- 10 11. A method according to any one of the claims 2-10, wherein said electrodes are positioned by use of a microscope, at least one micropositioner and/or a stereotactic device.

12. A method according to any one of the claims 2-11, wherein at least one electrode is used for delivery of at least one fusion partner to the fusion site.

13. A method according to any one of the claims 2-12, wherein step A is performed by use of the electrodes.

14. A method according to any one of the claims 1-12, wherein step A is performed by use of optical trapping.

15. A method according to any one of the claims 1-12, wherein step A is performed by use of micropipettes.

16. A method according to any one of the claims 1-15, wherein at least one of the fusion partners is a cell, and the other fusion partner(s) independently is (are) selected from the group consisting of a single cell, a liposome, a proteoliposome, a synthetic vesicle, an egg cell, an enucleated egg cell, a sperm cell at any development stage and a plant protoplast.

17. A method according to any one of the claims 1-15, wherein at least one of the fusion partners is constituted by a multiple of a structure selected from the group consisting of a single cell, a liposome, a proteoliposome, a synthetic vesicle, an egg cell, an enucleated egg cell, a sperm cell at any development stage and a plant protoplast.

18. A method according to any one of the claims 1-17, wherein the fusion partners are provided in a buffer prior to step B.

19. A method according to any one of the claims 1-18, wherein at least one of the fusion partners has been immobilized prior to step A.

20. A method according to any one of the claims 1-19, wherein one of the fusion partners is part of a cellular network.

21. A method according to any one of the claims 1-20, wherein at least one of the Fusion Partners has been electroporated in a buffer prior to step A.

22. A method according to any one of the claims 1-21, wherein at least one of the Fusion Partners has been exposed to a dielectrophoretic field in a buffer prior to step A.

23. A method according to any one of the claims 1-22, wherein at least one of the Fusion Partners has been treated by a fusogenic or other agent that promotes close cell-cell contacts.

24. Use of a method according to any one of the claims 1-23 in in vitro-fertilization.

25. Use of a method according to any one of the claims 1-23 in cloning.

26. Use of a method according to any one of the claims 1-23 for creation of hybridomas.

27. Use of a method according to any one of the claims 1-23 for manipulation of the composition of a cellular membrane.

28. Use of a method according to any one of the claims 1-23 for the delivery of a well-defined volume of a substance to a cell.

29. Use of a method according to any one of the claims 1-23 for the delivery of a pharmaceutically active substance to a cell.

30. Use of a method according to any one of the claims 1-23 in treatment of a tumor.

31. Use of a method according to any one of the claims 1-23 in treatment of chronic neurodegenerative diseases, such as Parkinson's disease and Alzheimer's disease.

---

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
2216  
2217  
2218  
2219  
2220  
2221


ABSTRACT

Disclosed is a method for selective electrofusion of  
at least two fusion partners having cell-like membranes  
5 and cellular or subcellular dimensions, comprising the  
following steps: A) the fusion partners are brought into  
contact with each other and B) an electrical field of a  
~~strength sufficient to obtain fusion and highly focused~~  
on the fusion partners is applied. The fusion partners  
10 are independently selected from the group consisting of a  
single cell, a liposome, a proteoliposome, a synthetic  
vesicle, an egg cell, an enucleated egg cell, a sperm  
cell at any development stage and a plant protoplast.


15

Publ. fig.: -


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100




**Fig. 1**



**Fig. 2**




**Fig. 3**



**Fig. 4**





**Fig. 5**

